

2008

Microbial Communities in Fire-Impacted Soils

T. C. Tobin
Susquehanna University

C. P. Janzen

Follow this and additional works at: http://scholarlycommons.susqu.edu/biol_fac_pubs



Part of the [Biology Commons](#)

Recommended Citation

Tobin, T. C., Janzen, C. P. (2008). Microbial Communities in Fire-Affected Soils. In Patrice Dion and Chandra Shekhar Nautiyal (Ed.), *Soil Biology Volume 13: Microbiology of Extreme Soils* (vol. 13, pp. 299-316). Berlin Heidelberg: Springer-Verlag.

This Book Chapter is brought to you for free and open access by Scholarly Commons. It has been accepted for inclusion in Biology Faculty Publications by an authorized administrator of Scholarly Commons. For more information, please contact sieczkiewicz@susqu.edu.

Chapter 13. Microbial Communities in Fire-Impacted Soils

1 Microbial Communities in Fire-Impacted Soils

1.1 Introduction

Of all the ways in which human activities can affect soil environments, fires are perhaps the most dramatic. Whether set deliberately or accidentally, the worldwide impact that such fires have is extensive, including the loss of human and animal lives as well as economic and ecological damage (UNECE et al. 2000; UNECE and FAO 2001; Davidenko and Eritsov 2003; Kudoh 2005; FAO 2005e). The United Nations Food and Agricultural Organization (2005) estimated that 350 million hectares burn annually, and that approximately 90% of those fires are of human origin. The frequency and severity of surface fires has also increased in many parts of the world due to changes in climate and land management practices (Houghton et al. 1992; Renkin and Despain 1992; Glantz 1996; Neary et al. 1999; Westerling et al. 2006). Large fires tend to draw media attention, particularly when they impinge on densely inhabited or well-known wildlife areas. Their rapid spread, extreme temperatures and the barren landscapes they leave behind can alter the surrounding ecosystem for years, decades, or even permanently. It is no surprise, therefore, that a premium has historically been placed on extinguishing fires as they happen, rather than on studying their ecological significance. Only recently have scientists begun to understand the critical roles that some fires play in sustaining natural environments, and on the parts that soil microorganisms play in that process.

Two basic types of fires will be discussed in this chapter, surface fires and underground fires. Surface fires include both wildfires and prescribed fires and their impacts are predominantly 'top down'. That is, the fire source is above ground, and the heat from the fire, while it can be intense enough to sterilize the surface soil, may not penetrate more than a few centimeters below the soil surface. Surface fire impacts often consist of a patchwork of severely affected sites interspersed with less affected areas, the pattern of which is dictated by the availability of fuel, topography, and other factors.

Underground fires, by way of contrast, have a 'bottom up' impact. They are typified by the large coal mine fires found in the Jharia Coalfield of India (Agarwal et al. 2006), throughout much of northern China (Zhou et al. 2006) and throughout the coal mining regions of the United States, most notably in Centralia, Pennsylvania (Trifonoff 2000). In these fires, steam and gases carrying vaporized combustion products from the burning coal below migrate, or 'vent', upward through soil fractures, cooling as they rise. As the gases from these 'anthracite smokers' cool, their dissolved chemicals either escape to the atmosphere or condense into the surrounding soils. Thus, underground fires are distinguished by subsurface soil temperatures that are generally much hotter than surface temperatures, and by soil chemical changes that tend to be

clustered around actively venting soil fractures. In many ways, these fires resemble geothermal environments as closely as they do environments impacted by surface fires. In this chapter, we will be using coalmine fires as the model system for underground fires in general.

Regardless of whether a fire has a top-down or a bottom-up impact, it alters many surface and subsurface soil properties as it migrates through an area. These properties include the availability of water, the structure and composition of soil particles, and the availability of nutrients. Sulfur, nitrogen, phosphorous and carbon species, in particular, can experience dramatic shifts when pre- and post-fire environments are compared.

The extent and severity of these fire-induced changes is determined by many factors including the duration of the fire, the intensity of the fire, the rate of the fire's spread, the topography of the burn area, the initial soil composition, the soil moisture content, fuel characteristics, weather conditions, the interval between fire events, and the frequency of fire events. However, the greatest of these is duration. A fast-moving above ground fire of even moderate intensity does not persist long enough to transfer enough heat to the soils to cause substantial belowground chemical changes. Slow-moving fires cause severe and complex changes that can be long lasting or permanent. Fire temperatures can range from 50° to >1500°C. Heat release can range from 2.11kJ/kg of fuel to 2.1MJ/kg. The effect of fire intensity is related to how well the energy produced by the fire is transferred to the soil. A slow moving intense fire will have a greater impact than a fast moving fire of the same intensity. Likewise, a slow moving fire of low intensity may have as great an impact as a fast moving fire due to the greater length of time the fire will be transferring the generated heat to the soil.

Ultimately, it is both the intensity and persistence of fires, as well as the resulting changes to the surface and subsurface soil properties that determine the fire's impact on resident microbial communities. Although these communities are likely to play critical roles in both post-fire biogeochemical nutrient cycling and in the ultimate recovery of fire-impacted environments, they have remained quite poorly studied. Most prokaryotic research, in particular, has focused on studies of post-fire microbial biomass, microbial metabolic assays, and culture-dependent assays (Ahlgren 1974; Dunn et al. 1979; Bissett and Parkinson 1980; Klopatek and Klopatek 1987; Klopatek et al. 1990; Fritze et al. 1993; Vazquez et al. 1993; Acea and Carballas 1996; Hernandez et al. 1997; Ross et al. 1997; Prieto-Fernández et al. 1998; Choromanska and DeLuca 2002; Andersson et al. 2004; Treseder et al. 2004; Naumova 2005; Giai and Boerner 2007). Only recently have culture-independent methods such as 16S rRNA gene sequencing, terminal restriction fragment length polymorphism (T-RFLP) analysis, phospholipids fatty acid analysis (PLFA) and denaturing gel gradient electrophoresis (DGGE) been combined with more traditional methods to give a more complete picture of the impacts that fires have on specific prokaryotic communities, and on the roles that those communities play in the post fire environment (Baath et al. 1995;

Pietikainen et al. 2000; Jaatinen et al. 2004; Tobin-Janzen et al. 2005; Yeager et al. 2005; Izzo et al. 2006).

1.2 The Impacts of Surface Fires on Microbial Environments

Since the fire-induced thermal, chemical and physical changes to the soil environment directly drive microbial community responses, it is important to understand these factors before turning to a more complete discussion of microbial ecology. The soil changes that most clearly impact microbial community responses include the intensity of the fire, and the resulting changes to the availability of water, nitrogen, sulfur, phosphorous and carbon.

1.2.1 Fire Intensity

The intensity of a surface fire, and its resulting impact on the microbial communities in underlying soils is due to the total amount of heat produced by the fire, the mode of transfer of that heat to the underlying soils, and the speed with which the fire progresses through an area. The total heat produced by the fire is governed by the nature of the fuel source and the fuel load. For example, grassland fires (Raison 1979; Ross et al. 1997; Neary et al. 1999) tend to have lower fuel loads and thus generate lower fire temperatures than fires in tropical woodlands or conifer forests. Since microbial mortality generally begins at around 50°C, but varies tremendously from species to species, the total heat produced by the fire is of paramount importance in determining the post-fire microbial community composition.

The modes of energy transfer from a surface fire to the soil are radiation, convection, conduction, vaporization/condensation, and mass transport (Neary et al. 1999). Electromagnetic radiation from the fire can play a significant role in energy transfer at the onset of a burn. Movement of air masses by convection can transfer heat from the active fire to nearby areas, heating nearby soils as well as spreading the fire. Conduction of heat by direct contact between hot or burning fuel and cooler soil can be important especially where fuel loads are heavy or the where fuels themselves are massive such as with slash piles. The high specific heat capacity of water ($4.184\text{J g}^{-1} \text{ }^{\circ}\text{C}^{-1}$) means that a substantial amount of energy is required to vaporize the water in soil. The enthalpies of heating and vaporization of soil moisture tend to mitigate any rise in soil temperature until all the water is vaporized. Indeed, the soil temperature cannot rise above 95°C until vaporization is complete (Campbell et al. 1994). The vaporized water carries latent heat through soils faster and deeper than other modes of energy transport such as radiation or conduction. Thus, while moist soils initially suffer lower temperatures, the overall impact of the fire may be more rapid and impact soil to a greater depth. The transfer of energy via mass transport generally has little to no effect on soil temperatures.

Once all of the moisture in soils under a fire have vaporized, soil temperatures typically rise to 200-300°C. However, under severe or slow moving fires, surface soil temperatures can rise to as high as 700°C (DeBano et al. 1998). The depth-temperature profile for a soil under a fire is determined by intensity and duration of the fire as well as the moisture content of the soil. With fast-moving or low severity fires, below ground soil temperatures generally do not exceed 100-150°C at 5 cm depth and demonstrate no heating below 30 cm (Agee 1973; DeBano 2000). A slow-moving or intense fire will clearly have a more significant effect on both the maximum soil temperatures observed and the depth at which temperatures are raised.

1.2.2 Available Moisture

Available soil moisture levels are not only decreased initially as a direct result of vaporization, but may also suffer long-term reductions. At moderate temperatures, incomplete burning of organic matter can result in the formation of a hydrophobic coating in the mineral components of soil resulting in increased repellency and decreased soil permeability. This repellency can result in a persistent decrease in soil moisture in the post burn soils or in greater run-off and erosion. Despite these tendencies, situations in which soil moisture has increased (Klock and Hevey 1976; Haase 1986) or remained the same (Campbell et al. 1977; Milne 1979) after fire have also been documented, and thus the overall impact that post-fire soil moisture has on microbial communities is varied (Letey 2001).

1.2.3 Available Nutrients

In an above ground fire, the most dramatic and well-understood changes to available nutrients involve the volatilization, chemical transformation and biogeochemical cycling of nitrogen, carbon, phosphorous and sulfur. While changes can and do occur to trace minerals such as As, Ca, K, Na, Fe and Al (Grier 1975; Feller 1982; Macadam 1989; DeBano et al. 1998; Neary et al. 1999; Arocena and Opio 2003)) the impacts that these changes have on microbial communities remain mostly unstudied, and thus will not be discussed in this chapter.

Nitrogen. At moderate temperatures, dead partially combusted plant and microbial biomass can be easily oxidized resulting in an increase in inorganic nitrogen immediately post burn (Diaz-Ravina et al. 1996). More extreme temperatures can result in volatilization of nitrogen (Giovannini et al. 1990). Total nitrogen decreases slightly as the temperature rises from 25 to 220°C but dramatically decreases between 220 and 460°C (Giovannini et al. 1990). Ammonium concentrations steadily increase up to 220°C but, like the total nitrogen, drop quickly at the temperature rises above 220°C until very little remains at 460°C. It is presumed that this increase in NH₄-N is due to mineralization of organic nitrogen. Nitrate concentrations are initially unaffected

by fire, even moderate to intense fires. However, NO_3^- - N is produced by biochemical nitrification of ammonium in the time following the fire resulting in NO_3^- - N concentrations that can be significantly higher in the weeks or years post-fire (Covington et al. 1991). Therefore, while total nitrogen may decrease, the bioavailable forms of nitrogen, nitrate and ammonium, may both be at elevated levels for several years following a burn. Ammonium tends to adsorb onto the mineral soil and become immobilized. Unless regrowth of vegetation occurs soon after the fire, the nitrate can be easily lost to leaching, thus depleting the total nitrogen for long periods. Conversely, if regrowth is rapid, soil organic nitrogen concentrations can rapidly recover to pre-fire levels (Adams and Attiwill 1984; Weston and Attiwill 1996).

Carbon. Fire severity as described by the maximum sustained surface temperature under a fire affects the degree of consumption of litter and soil organic matter. A low severity burn where the soil temperatures do not exceed 250°C can cause partial scorching of litter. As depth in the soil increases, the insulating ability of the soil mitigates the heating of soils. Thus with surface temperatures at or below 250°C , soil temperatures down to 2.5 cm rarely exceed 100°C and at 5 cm the temperature is typically below 50°C . The lower temperatures at depth result in only partial distillation of organic matter above 2.5 cm and little to no effect below 5 cm. A moderate burn characterized by surface temperatures up to 400°C results in temperatures at 2.5 and 5 cm of up to 175 and 50°C , respectively. These temperatures result in very significant charring of litter, some charring of organic matter to 2.5 cm and the start of distillation of organic matter above 5 cm. In a severe burn surface temperatures exceed 675°C , resulting in complete combustion of the litter. The soil down to 2.5 cm can experience heating to 190°C with concomitant charring or consumption of large portions of the organic matter. Even at depths of 5 cm soils are being heated to 75°C resulting in significant distillation of volatile organic matter and some charring (DeBano et al. 1977).

Phosphorous. Organic phosphorous is readily converted to inorganic forms of phosphate. Even in low severity fires where temperatures do not exceed 200°C , the concentration of organic phosphorous decreases markedly. In an artificial heating experiment, organic phosphorous was completely depleted from soils above 220°C (Giovannini et al. 1990). While organic carbon is easily lost, total phosphorous remains fairly constant due to its low volatility. The organic phosphate is instead converted to orthophosphate, the predominate form of bioavailable phosphorous (Cade-Menun et al. 2000). The concentrations of inorganic and available phosphorous peak at soil temperatures of approximately 450°C (Giovannini et al. 1990). Other factors play a role in determining how much of the inorganic phosphorous produced by a burn is available. Bioavailability peaks at a pH around 6.5 (Sharpley 2000). The ash produced in a fire tends to move the soil pH higher toward this value. However, in calcareous soils phosphate can complex very strongly to calcium resulting in removal of phosphorous from the available pool. In acidic soils, phosphate can bind with

other soil metals such as iron, manganese, and aluminum. In summation, above-ground fires easily convert most or all of the organic phosphorous to inorganic phosphorous, thus initially increasing the bioavailability. How long the increase in bioavailable phosphorous persists depends on soil pH, the composition of the mineral soil, and the rate of uptake of available phosphorous by recolonizing vegetation.

Sulfur. In an artificial heating study, Badia and Marti (Badia and Marti 2003) found that a slight steady increase in total sulfur occurred upon heating gypsiferous soils up to 500°C and upon incorporation of ash into the soil, but that no change in total sulfur occurred in calcerous soils. In another study, Castelli and Lazzari (Castelli and Lazzari 2002) found that total sulfur peaked a year after a controlled burn in both grass-covered and shrub-covered soils. After a second controlled burn three years after the first burn, the total sulfur in the soils under grass cover dropped back to the original (before the first burn) levels below 1 cm depths. In the same study, available sulfur under shrub cover immediately jumped significantly, but then dropped back to pre-burn levels within two years. The available sulfur under grass cover was not affected. In both cases, the available sulfur did not change with the second controlled burn. These two studies indicate that the sulfur chemistry of soils is impacted by above ground fires in manners similar to other nutrients.

1.3 The Impacts of Underground Fires on Microbial Environments

Underground coal mine fires have very different effects on soil chemistry than those caused by aboveground fires. The reasons stem from three major factors. The heat source is below the surface, the duration of the belowground fires can be considerably longer than surface fires, and the composition of the coal fuel is quite different than that found in typical prescribed or wild aboveground fires. These factors lead to unique physical and chemical properties and processes.

1.3.1 Fire Intensity

In an aboveground fire, the highest temperatures the soils experience are at the air-surface interface. Conversely, the lowest temperature of an affected soil is at the air-surface interface with a belowground fire. In other words, the soil temperature gradient in a belowground fire is inverted from that observed in aboveground fires.

In addition, the impact of the fire is not evenly distributed spatially. The primary modes of energy transfer from the fire to the surface and near-surface soils are convection and conduction. Convection of hot combustion gases takes the paths of least resistance, following the small fractures and faults in the subsurface. The gases tend to escape to the atmosphere in vents. The temperature of the surrounding soils is highest near the vent as a result of the transfer of thermal energy from the hot gases to the surrounding soils. These hot soils are then able

to transfer heat away from the vent via conduction. Eventually, this heat is slowly lost to the atmosphere. Thus the surface temperature decreases with increasing distance from the vent. In addition to heat brought to the surface by convection, the hot subsurface rock and mineral matter can transfer heat to the surface directly by conduction. This conduction results in surface regions that are quite hot while displaying no outward signs of venting of hot gases. Indeed, the authors have measured surface temperatures above the Centralia Pennsylvania mine fire exceeding 400°C in non-venting areas (Tobin-Janzen unpublished results). The underground environment (including both structure and composition) is not uniform. Therefore, the heat transferred by conduction is spatially uneven. The end result of the modes of heat transfer is that the surface temperatures above an underground fire are distributed in an irregular pattern.

Another significant difference between aboveground and belowground fires is the duration. Aboveground fires persist at a certain location for hours to weeks. The available fuel is consumed quickly and when gone, no further burning is possible. Stated another way, aboveground fires are fuel limited. The rate of progression of belowground fires is much slower, and can range from ten meters per year to several hundred meters per year. Even at the fastest rates, underground fires progress more slowly than aboveground fires. This slow movement means that an area on the surface that becomes impacted by the underground fire may experience these effects for years. Indeed, certain areas in Centralia have been affected continuously for decades (Trifonoff 2000).

The principal reason for the slow movement is a shortage of oxygen. That is, belowground fires are oxidant limited. Oxygen is replenished through the fractures in the subsurface structures. Indeed, weather systems affect the progress of the fire. Low-pressure weather systems result in more venting of gases whereas little venting takes place during high-pressure systems. Another consequence of this shortage of oxygen is that the underground fire is a reducing fire. Thus, the reduced and volatile forms of nitrogen (ammonium) and sulfur (S° and H_2S) are formed and transported to the surface.

1.3.2 Available Moisture

In general, as the hot combustion products of the fire rise through the mineral and soil column those components with low volatilities begin to condense. One simple manifestation of this phenomenon is the formation of a very moist semi-liquid layer beneath active vents. When the hot gas encounters cooler soil, water condenses creating a very hot mud between 0.5 and 1m beneath the surface. (Tobin-Janzen unpublished results). Above this level, the soil moisture varies tremendously, and can even be quite dry, more closely resembling soils impacted by above ground heat sources (Tobin-Janzen et al. 2005).

1.3.3 Available Nutrients

The chemical composition of coal is markedly different from the organic fuel burned in aboveground fires. Coal is a highly variable fossilized form of ancient plant matter that has been significantly altered by exposure to elevated temperature and pressure. In addition to the organic components, coal contains significant inorganic materials. When burned, the inorganic components can melt and volatilize. Indeed, these inorganic phases pose technological problems for the use of coal as a clean energy source. The combustion products, including the inorganic phases, can percolate to the surface and near-surface environments, where they have been shown to condense into a variety of minerals, including elemental sulfur, downeyite (SeO_2), orpiment (As_2S_3), laphamite ($\text{As}_2(\text{Se},\text{S})_3$), ammonium chloride, gypsum ($\text{CaSO}_4\cdot\text{H}_2\text{O}$), and mullite ($\text{Al}_6\text{Si}_2\text{O}_{13}$) (Finkelman and Mrose 1977; Lapham et al. 1980; Dunn et al. 1986). Indeed, the authors have employed X-ray powder diffraction to identify pure ammonium chloride and needle-like elemental sulfur crystallizing on the surface of the ground above the Centralia Pennsylvania mine fire (Janzen, unpublished results). While the rate of deposition of these compounds is not high, the impact caused by a long and slow-burning underground coalmine fire has the potential to be significant. Also, as described above, the distribution of combustion gases and volatile chemical species at the surface is not even. Rather, it is a function of the sub-surface structure. Thus localized areas of high chemical concentrations, temperature, and soil moisture are interspersed with areas of lower concentrations and more moderate temperatures.

Nitrogen Tobin-Janzen et al (2005) studied the inorganic nitrogen ($\text{NH}_4\text{-N}$ and $\text{NO}_3\text{-N}$) levels in surface soils overlying the Centralia, Pennsylvania coal mine fire over a two-year period. During that time, the mine fire front progressed through the sample site, and surface soil temperatures increased an average of 16.1°C . While many of the hottest areas showed elevated ammonium (as high as 13.79 mg/kg) and nitrate (as high as 103.1 mg/kg) levels, these concentrations were not correlated with absolute temperature values, nor to proximity to an active vent, as some areas with elevated temperature, close proximity to an active vent, or both also had inorganic nitrogen concentrations that were comparable to those in nearby unaffected soils. Rather, inorganic nitrogen levels were related to more complex environmental trends, such as the length of time an area had been impacted by the mine fire and whether the overall soil temperatures were rising or falling in the area due to the movement of the mine fire front.

Sulfur The sulfur containing gases percolating up from the burning coal have a profound effect on the soil chemistry of the overlying soils, and thus represent one of the most striking differences between underground mine fire-impacted soils and those impacted by surface fires. For example, in unpublished work, the total sulfur concentrations are clearly impacted by the underground fire (Fig. 1). The concentrations are significantly higher near the vent. Interestingly, the sulfur concentrations are not highest directly at the vent. Rather, they increase to a maximum approximately 0.5m from the vent, then decrease with distance from

the vent. At greater distances from the main vent the concentrations of total sulfur again begin to drop off due to the depletion of volatile sulfur components of the gases at greater distances from the source. Similar results have been observed for elemental sulfur (S^0).

Using thermodynamic principles, Stracher has proposed a stability relationship between the partial pressure of S_2 gas and temperature (Stracher 1995). He developed a model predicting which phase, orthorhombic or S_2 gas, would be stable under a given set of conditions. Depending on the partial pressure of S_2 the temperature at which the solid form of elemental sulfur becomes the favored species varies. At very low partial pressures of S_2 , sulfur prefers to be solid at low temperatures. As the pressure rises, the solid form becomes favored at higher temperatures. Thus, at the vent where temperatures are highest, the gaseous form is predominant. Moving a short distance from the vent, the temperature is lower, allowing sulfur to solidify. Further from the vent, the sulfur has already been depleted in the effluent gas, thus no further deposition is possible.

1.4 Microbial Communities in Fire-Impacted Soils

The response of microbial communities to fire varies primarily with the nature of the fire. That is, whether the fire has a top-down or a bottom-up profile, its intensity and duration, and the resulting chemical, physical and biological changes it generates in the surrounding environment. Not surprisingly, the fire's intensity and persistence tend to be the most important factors in governing post-fire microbial community numbers and diversity. Nevertheless, other factors, such as the release of organic toxins, including polychlorinated dibenzo – p-dioxins (PCDDs), dibenzofurans (PCDFs), and polynuclear aromatic hydrocarbons (PAHs), into the surrounding soils can dramatically and negatively impact microbial biomass and diversity (Kim et al. 2003). Conversely, the increase of inorganic nitrogen and sulfur levels in fire-impacted soils can allow microbes capable of exploiting these molecules in their metabolic strategies to thrive in many post-fire environments, particularly if they are able to withstand the fire's high temperatures or to re-colonize fire-impacted areas quickly as they cool.

1.4.1 Fire Intensity and Frequency

The fire's intensity, as determined by its total heat, the transfer of that heat to the under- or overlying soils, and the migration rate of the fire is generally the most important single factor driving microbial community responses in most environments (Neary et al. 1999; Hart et al. 2005; Tobin-Janzen et al. 2005). Heat from the fire is by itself often high enough to sterilize soils, with surface temperatures above mine fires exceeding 730°C (Lapham et al. 1980), and those in surface fires capable of exceeding 675°C . Even non-sterilizing heat levels can directly lyse bacterial cells, and reduce their reproductive capabilities (Covington and DeBano 1990). As a result, total microbial numbers as well as microbial

community diversity tend to decrease dramatically immediately post-fire (Dunn et al. 1979; Fritze et al. 1993; Prieto-Fernández et al. 1998; Tobin-Janzen et al. 2005), although some exceptions to this rule have been observed (Newman et al. 2003).

Soil moisture, which varies with both above and below ground fire conditions as previously described, can intensify the fire's impact on microbial cells. Moist heat is much more efficient at killing soil microorganisms than dry heat, with threshold values for fungal and bacterial survival estimated at 80°C and 60°C for fungi, and 120°C and 100°C for bacteria in dry and moist soils, respectively (Dunn and DeBano 1977; Dunn et al. 1985). Climate can also impact soil moisture. Drought conditions in the summers following both surface and underground fires have been proposed to be involved in the post-fire persistence of decreased bacterial and fungal biomass (Cilliers et al. 2005; Tobin-Janzen et al. 2005; Yeager et al. 2005), although the condensing steam from underground mine fires may actually help to ameliorate drought conditions in some instances (Tobin-Janzen et al. 2005).

The migration rate of a fire likewise determines its impact on soil microbial communities. Low intensity, rapidly moving surface fires do not generally exceed surface temperatures of 250°C, and microbes at depths as shallow as 25 mm generally survive. By contrast, in high intensity surface fires with surface temperatures exceeding 675°C, microbes as deep as 50 mm underground experience selective die-off (Neary et al. 1999). By comparison to surface fires, underground fires tend to move very slowly, and their impacts can last for decades or longer. The Centralia mine fire, which started in 1962, is expected to burn for at least another 100 years, and the Jharia coalmine fire has been burning in India since 1916. Thus, the long-term impacts to soil microbial communities in these areas, while yet unstudied, are expected to be extensive. Over a one-year period, our lab (Tobin-Janzen et al. 2005) used T-RFLP analysis of 16S rRNA domain Bacteria ribotypes to ascertain that the complexity of microbial communities decreased uniformly as temperatures in the site increased. Furthermore, these changes were not correlated to other environmental factors such as soil pH, soil moisture, inorganic nitrogen or total sulfur, but rather with general trends in the mine fire progression itself. It will be of interest to see what happens to these microbial communities as the impact time stretches from months to decades.

Another important manner in which the intensity of surface fires and underground fires differ from one another is in the bottom-up nature of underground fires. Since the fire source is below ground, subsurface temperatures become progressively more elevated as soil depth increases. Although the subsurface microbial communities below 50 cm have not yet been extensively assayed in our lab, our unpublished observations suggest that as temperatures increase with depth, the total microbial biomass, as determined by the amount of bacterial DNA that can be extracted from the soil samples, decreases as well. Despite

this decrease, we have successfully amplified domain Bacteria 16S rRNA genes from soil samples at depths of up to 50 cm, and at temperatures of up to 87°C.

The overall impact of a fire's intensity on microbial communities varies tremendously with the species composition of that community. Pietikainen et al. (2000) demonstrated that bacteria tend to be more resistant to fire-induced heat than fungi. However, not all bacteria are similarly resistant, nor are all fungi similarly susceptible. In fact, Izzo et al. (2006) used greenhouse experiments to demonstrate that certain species of ectomycorrhizal fungi may actually compete more successfully for roots in the presence of fire, and *Neurospora* ascospores germinate in response to heat or fire (Emerson 1948; Jesenska et al. 1993; Pandit and Maheshwari 1996).

Heterotrophic bacteria often show the sharpest declines in numbers following fires, presumably as a result of the combustion of readily available food sources. By contrast, autotrophic bacterial numbers can actually increase as nutrients are released from their organic forms. In our work above the Centralia coalmine fire, for example, autotrophs are frequently dominant members of the bacterial communities present in fire-impacted soils. However, *Nitrobacter* are generally more heat sensitive than heterotrophic bacteria, (Dunn and DeBano 1977; Dunn et al. 1985), and actinomycetes are generally more heat resilient than other culturable heterotrophs (Cilliers et al. 2005). Recent experiments have shown that many of the most common bacteria in fire-impacted soils include spore formers (Ahlgren 1974; Moseby et al. 2000; Yeager et al. 2005). This finding is not surprising as these spore formers probably survive short-term intense heat as spores, and then rapidly vegetate once soil moisture, temperature and nutrient levels have returned to sustaining levels. Ultimately, thermophilic bacteria are the most resilient of all microbial species, and are often preferentially selected for by fire conditions. Our lab and others have demonstrated that they often make up the dominant populations in fire-impacted soils (Norris et al. 2002; Tobin-Janzen et al. 2005; Yeager et al. 2005).

1.4.2 Nitrogen-Cycling Bacteria

Available nitrogen is often a limiting nutrient in pre-fire environments (Allen et al. 2002; Yeager et al. 2005). However, both surface and underground fires can release inorganic nitrogen as a result of the combustion of organic molecules. This nitrogen initially exists as $\text{NH}_4\text{-N}$, but is quickly oxidized to $\text{NO}_3\text{-N}$ (DeLuca and Zouhar 2000; Tobin-Janzen et al. 2005; Yeager et al. 2005). Soil bacteria are probably critical components in this oxidation, and yet they have remained poorly studied in post-fire environments until recently. Yeager et al. (2005) used molecular analysis of *nifH* and *amoA* genes to study nitrogen-fixing and nitrifying bacteria following a forest fire. They demonstrated that while there was a decrease in overall microbial biomass, including that of nitrogen cycling bacteria, the nitrogen-fixing community actually became more diverse within a month after the fire. By contrast, a single ammonia-oxidizing type, belonging to cluster 3A

Nitrosospira spp., dominated in the post-fire soils. Our lab has similarly demonstrated the presence of ammonia-oxidizing bacteria, as determined by the presence of *amoA* genes, in hot mine-fire impacted soils with elevated $\text{NH}_4\text{-N}$ and $\text{NO}_3\text{-N}$, and at temperatures of up to 60°C (Tobin-Janzen et al. 2005). Furthermore, we have isolated spore-forming *Geobacilli* that can reduce nitrate to N_2O at 60°C (Kauffman and Tobin-Janzen 2005). Thus, nitrogen cycling bacteria appear to be important components of both surface and underground mine fire environments, where they are most likely exploiting the elevated nitrogen levels, and where they could be contributing to greenhouse emissions in the fire areas.

1.4.3 Sulfur-Cycling Bacteria

Perhaps the most interesting difference between environments impacted by belowground versus surface fires is the high levels of sulfur that can condense into the soils above the former. In this regard, subsurface fires tend to more closely resemble fumaroles and other similar geothermal environments, in which surface soils are impacted by hot gases containing varying levels of H_2O , CO_2 , SO_2 , H_2S , H_2 , N_2 , CO and CH_4 (A chemical study of Hawaiian magmatic gases, Geochemistry of the magmatic-hydrothermal system of Kawah Ijen Volcano, East Java, Indonesia; Variations in the temperature and chemistry of White Island fumarole discharges 1972-85).

Janzen, et al (unpublished) have recently studied the sulfur levels surrounding an active vent above the Centralia coalmine fire, where they demonstrated that the highest levels of sulfur did not necessarily correlate with the highest soil temperatures, but rather with temperatures that favored condensation of the sulfur into the soils, rather than its loss to vaporization (Fig.1). In order to determine if sulfur-metabolizing bacteria were capitalizing on the high levels of sulfur in these soils, our lab collected sterile soil samples from the same boreholes, and identified total Bacterial community members using 16S rRNA gene sequencing. In Figure 2, it can be seen that the bacterial community members of this fire-impacted soil are in fact enriched for thermophilic bacteria, including thermophilic sulfur metabolizers, nitrogen cycling bacteria, and spore-forming bacteria. The community itself was only modestly diverse, with the dominant bacterial groups including *Actinobacteria*, *Alphaproteobacteria*, *Deltaproteobacteria*, *Betaproteobacteria*, along with *Acidothermus*, *Clostridium* and *Geobacillus* species. These results are in good agreement with the trends seen in other hot, fire- and geothermally-impacted soil environments (Norris et al. 2002; Yeager et al. 2005).

1.5 Conclusions

Although the microorganisms capable of surviving and thriving in fire-impacted areas are poorly studied at this time, studying their life cycles and environmental impacts is likely to produce important discoveries. Microbes found in fire-impacted areas most likely play critical roles in the biogeochemical cycling of

nitrogen and in rhizosphere ecology, and thus can be expected to play important roles in the recovery of fire-impacted ecosystems. Thermophilic bacteria, in general, have already provided biotechnology with some of its most powerful tools, and thermophilic sulfur-metabolizing bacteria are currently at the heart of several studies geared toward producing more efficient methods of bioleaching metals and desulfurizing flue gases (Huber and Stetter 1998; Kaksonen et al. 2006). Since these bacteria appear to be enriched in hot mine-fire impacted soils, continued research into their biology, environmental roles and metabolism will be particularly interesting. Finally, not all fire-associated bacteria are benign. Iron and sulfur-metabolizing bacteria catalyze the rate limiting step responsible for acid mine drainage in coal mining environments, and the nitrogen cycling bacteria identified in this study and others may also be responsible for releasing the greenhouse gases NO and N₂O into the environment. Thus, the more we learn about these interesting extremophiles, the more we are likely to benefit from that knowledge.

Figure Legends

Figure 13.1 Spatial distribution of total sulfur around an active vent above the Centralia, Pennsylvania mine fire. The concentrations of total sulfur were measured in dry soil on two days during the summer of 2005. The results are correlated with soil temperature and distance from the vent. Triplicate analyses were performed on each sample and the error bars represent the 95% confidence intervals.





Figure 13.2 Dendrogram of 16S rRNA gene sequences obtained from the field samples surrounding the active vent depicted in figure 13.1. Samples designated 'Clone C' and 'Clone I' originated from the boreholes one meter to the left and right of the active vent, respectively. Samples designated Clone A or Clone U came from an affected surface site (50°C) located outside of the study area depicted in Figure 13.1. Taxa containing known thermophiles (T), nitrogen (N), sulfur (S) and iron (Fe) metabolizers are indicated. The Dendrogram was generated using Nearest Neighbor analysis and the CLC Combined Workbench Program (CLC Bio) Bootstrap values from 100 resamplings are shown above each internal node. *Thermaerobacter* 16S rRNA was used as the out-group



References Cited

Acea MJ and Carballas T (1996) Changes in physiological groups of microorganisms in soil following wildfire. *FEMS Microbiology Ecology* 20:33-39

Adams MA and Attiwill PM (1984) Role of *Acacia* spp. in nutrient balance and cycling in regenerating *Eucalyptus regnans* F. Muell. Forests. I. Temporal changes in biomass and nutrient content. *Australian Journal of Botany* 32:205-215

Agarwal R, Singh D, Chauhan DS and Singh KP (2006) Detection of coal mine fires in the Jharia coal field using NOAA/AVHRR data. *Journal of Geophysics and Engineering* 3:212-218

Agee JK (1973) Prescribed Fire Effects on Physical and Hydrologic Properties of Mixed-conifer Forest Floor and Soil. Report 143 University of California Resources Center 143.

Ahlgren IF (1974) The effect of fire on soil organisms. In: Kozlowski, T.T. and Ahlgren, C.E., eds) *Fire and ecosystems*. Academic press, New York, pp47-72

Allen CD, Savage M, Falk DA, Suckling KF, Swetnam TW, Schulke T, Stacey PB, Morgan P, Hoffman M and Klingel JT (2002) Ecological restoration of Southwestern ponderosa pine ecosystems: a broad perspective. *Ecological Applications* 12:1418-1433

Andersson M, Michelsen A, Jensen M and Kjoller A (2004) Tropical savannah woodland: effects of experimental fire on soil microorganisms and soil emissions of carbon dioxide. *Soil Biology and Biochemistry* 36:849-858

Arocena JM and Opio C (2003) Prescribed fire-induced changes in properties of sub-boreal forest soils. *Geoderma* 113:1-16

Baath E, Frostegard A, Pennanen T and Fritze H (1995) Microbial community structure and pH response in relation to soil organic matter quality in wood-ash fertilized, clear-cut or burned coniferous forest soils. *Soil Biology and Biochemistry* 27:229-240

Badia D and Marti C (2003) Plant Ash and Heat Intensity Effects on Chemical and Physical Properties of Two Contrasting Soils. *Arid Land Research and Management* 17:23-41

Bissett J and Parkinson D (1980) Long-term effects of fire on the composition and activity of the soil microflora of a subalpine, coniferous forest. *Canadian Journal of Botany* 58:1704-1721

Cade-Menun BJ, Berch SM, Preston CM and Lavkulich LM (2000) Phosphorous forms and related soil chemistry of Podzolic soils on northern Vancouver Island. II. The effects of clear-cutting and burning. *Canadian journal of forest Research* 30:1726-1741

Campbell GS, Jungbauer JD, Bidlake WR and Hungerford RD (1994) Predicting the effect of temperature on soil thermal conductivity. *Soil Science* 158:307-313

Campbell RE, Baker MB, Ffolliott PF, Larson FR and Avery CC (1977) Wildfire effects on a ponderosa pine ecosystem: an Arizona case study USDA For. Serv. Res. Pap. RM-191.

Castelli LM and Lazzari MA (2002) Impact of Fire on Soil Nutrients in Central Semiarid Argentina. *Arid Land Research and Management* 16:349-364

Choromanska U and DeLuca TH (2002) Microbial activity and nitrogen mineralization in forest mineral soils following heating: evaluation of post-fire effects. *Soil Biology and Biochemistry* 34:263-271

Cilliers CD, Botha A, Esler KJ and Boucher C (2005) Effects of alien plant management, fire and soil chemistry on selected soil microbial populations in the Table Mountain National Park, South Africa. *South African Journal of Botany* 71:211-220

Covington WW and DeBano LF (1990) Effects of fire on pinyon-juniper soils. In: Krammes, J.S., (ed. Effects of Fire Management of Southwestern Natural Resources. USDA For. Serv. Gen. Tech. Rept. RM-191, pp78-86

Covington WW, DeBano LF and Huntsberger TG (1991) Soil nitrogen changes associated with slash pile burning in pinyon-juniper woodlands. *Forest Science* 37:347-355

Davidenko EP and Eritsov A (2003) The fire season in 2002 in Russia. Report of the Aerial Forest Fire Service, Avialesookhrana. *International Forest Fire News* 28:15-17

DeBano LF (2000) The Role of Fire and Soil Heating on Water Repellency in Wildland Environments: A Review. *Journal of Hydrology* 231:195-206

DeBano LF, Dunn PH and Conrad CE (1977) Fire's effect on physical and chemical properties of chaparral soils. Mooney, H.A. and Conrad, C.E. (eds.) *Proceedings of the Symposium on the Environmental Consequences of Fire and Fuel Management in Mediterranean Ecosystems, Palo Alto, CA., USDA For. Serv. Gen. Tech. Rep. WO-3. pp65-74*

DeBano LF, Neary DG and Ffolliott PF (1998). *Fire Effects on Ecosystems*. Wiley, New York

DeLuca TH and Zouhar KL (2000) Effects of selection harvest and prescribed fire on the soil nitrogen status of ponderosa pine forests. *Forest Ecology and Management* 138:263-271

Diaz-Ravina M, Prieto A and Baath E (1996) Bacterial activity in a forest soil after heating and organic amendments measured by the thymidine and leucine incorporation technique. *Soil Biology and Biochemistry* 28:419-426

Dunn PH and DeBano LF (1977) Fire's effects on biological and chemical properties of chaparral soils. Mooney, H.A. and Conrad, C.E. eds.) *Proceedings of the Symposium on the Environmental Consequences of Fire and Fuel Management in Mediterranean Ecosystems, Palo Alto, CA., USDA For. Serv. Gen. Tech. Rep. WO-3. pp75-84*

Dunn PH, DeBano LF and Eberlein GE (1979) Effects of burning on chaparral soils: II Soil microbes and nitrogen mineralization. *Soil Science Society of America Journal* 43:509-514

Dunn PH, Barro SC and Poth M (1985) Soil moisture affects survival of microorganisms in heated chaparral soil. *Soil Biology and Biochemistry* 17:143-148

Dunn PJ, Peacor DR, Criddle AJ and Finkelman RB (1986) Laphamite, an arsenic selenide analogue of orpiment, from burning anthracite deposits in Pennsylvania. *Mineralogical Magazine* 50:279-282

Emerson M (1948) Chemical activation of ascospore germination in *Neurospora crassa*. *Journal of Bacteriology* 55:327-330

FAO (2005e) State of the World's Forests 2005

Feller MC (1982) The ecological effects of slash burning with particular reference to British Columbia: a literature review Province of British Columbia Ministry of Forests Land Management Report No. 13.

Finkelman RB and Mrose ME (1977) Downeyite, the first verified natural occurrence of SeO_2 . *American Mineralogist* 62:316-320

Fritze H, Pennanen T and Pietikäinen J (1993) Recovery of soil microbial biomass and activity from prescribed burning. *Canadian Journal of Forest Research* 23:1286-1290

Giai C and Boerner REJ (2007) Effects of ecological restoration on microbial activity, microbial functional diversity, and soil organic matter in mixed-oak forests of southern Ohio, USA. *Applied Soil Ecology* 35:281-290

Giovannini G, Lucchesi S and Giachetti M (1990) Effects of Heating on some Chemical Parameters Related to Soil Fertility and Plant Growth. *Soil Science* 149:344-350

Glantz MH (1996). In: *Currents of Change: El Niño's Impact on Climate and Society*. Cambridge University Press, Cambridge, p194

Grier CC (1975) Wildfire effects on nutrient distribution and leaching in a coniferous ecosystem. *Canadian Journal of Forest Research* 5:599-607

Haase SM (1986) Effect of prescribed burning on soil moisture and germination of southwestern ponderosa pine seed on basaltic soils USDA For. Serv., Rocky Mt. For. Range Exp. Stn. Res. Note RM-462.

Hart SC, DeLuca TH, Newman GS, MacKenzie MD and Boyle SI (2005) Post-fire vegetative dynamics as drivers of microbial community structure and function in forest soils. *Forest Ecology and Management Forest Soils Research: Theory, Reality and its Role in Technology - Selected and Edited Papers from the 10th North American Forest Soils Conference held in Saulte Ste. Marie, Ontario, Canada, 20-24 July 2003* 220:166-184

Hernandez T, Garcia C and Reinhardt I (1997) Short-term effect of wildfire on the chemical, biochemical and microbiological properties of Mediterranean pine forest soils. *Biology and Fertility of Soils* 25:109-116

Houghton JH, Callander BA and Varney SK (1992) *Climate Change 1992: The Supplementary Report to the IPCC Scientific Assessment* Press Syndicate of the University of Cambridge p220

Huber H and Stetter KO (1998) Hyperthermophiles and their possible potential in biotechnology. *Journal of Biotechnology* 64:39-52

Izzo A, Canright M and Bruns TD (2006) The effects of heat treatments on ectomycorrhizal resistant propagules and their ability to colonize bioassay seedlings. *Mycological Research* 110:196-202

Jaatinen K, Knief C, Dunfield PF, Yrjala K and Fritze H (2004) Methanotrophic bacteria in boreal forest soil after fire. *FEMS Microbiology Ecology* 50:195-202

Jesenska Z, Pieckova E and Bernat D (1993) Heat resistance of fungi from soil. *International Journal of Food Microbiology* 19:187-192

Kaksonen AH, Plumb JJ, Robertson WJ, Spring S, Schumann P, Franzmann PD and Puhakka JA (2006) Novel Thermophilic Sulfate-Reducing Bacteria from a Geothermally Active Underground Mine in Japan. *Applied and Environmental Microbiology* 72:3759-3762

Kauffman C and Tobin-Janzen T (2005) Isolation and Identification of Soil Microbial Populations in Centralia, Pennsylvania. *Journal of the Pennsylvania Academy of Sciences* 78:117

Kim E-J, Oh J-E and Chang Y-S (2003) Effects of forest fire on the level and distribution of PCDD/Fs and PAHs in soil. *The Science of The Total Environment* 311:177-189

Klock GO and Hevey JD (1976) Soil-water trends following wildfire on the Entiat Experimental Forest. *Ann. Proc. Tall Timber Fire Ecol. Conf.* 15:193-200

Klopatek CC and Klopatek JM (1987) Mycorrhizae, microbes, and nutrient cycling processes in pinyon-juniper systems. Everett, R.L. (compiler). *Proceedings of the Pinyon-Juniper Conference, January 13-16, 1986 Reno, NV, USDA For. Serv. Gen. Tech. Rep. INT-215. pp360-367*

Klopatek CC, DeBano LF and Klopatek JM (1990) Impact of fire on the microbial processes in pinyon-juniper woodlands: management implications. In: *Effects of Fire Management of Southwestern Natural Resources. USDA For. Serv. Gen. Tech. Rep. RM-191 pp197-205*

Kudoh J (2005) Report of the View of Northeast Asia Forest Fire from Cosmos International Symposium Center for Northeast Asian Studies (CNEAS), Tohoku University, Sendai, Miyagi, Japan, 17-18 January 2005, 17-18 January 2005.

Lapham DM, Barnes JH, Downey WF and Finkelman RB (1980) Minerology associated with burning anthracite deposits of eastern Pennsylvania Pennsylvania Geological Survey Mineral Research Report M 78.

Letey J (2001) Causes and consequences of fire-induced soil water repellency. *Hydrological Processes* 15:2867-2875

Macadam A (1989) Effects of prescribed fire on forest soils. BC Ministry of Forests Research Report 89001-PR, Victoria.

Milne MM (1979) The effects of burning root trenching and shading on mineral soil nutrients in southwestern ponderosa pine. M.S. Thesis Northern Arizona University, Flagstaff, AZ.

Moseby AH, Burgos J, Reed J and Tobin-Janzen T (2000) Isolation and Identification of Soil Bacteria from the Centralia Mine Fire Area. *Journal of the Pennsylvania Academy of Sciences* 73:150

Naumova NB (2005) Biomass and Activity of Soil Microorganisms after a Surface Fire in a Pine Forest. *Eurasian Soil Science* 38:870-873

Nearby DG, Klopatek CC, DeBano LF and Ffolliott PF (1999) Fire effects on belowground sustainability: a review and synthesis. *Forest Ecology and Management* 122:51-71

Newman GS, Hart SC, Guido D and Overby S (2003) Wildfire effects on soil microbial activity and community-level physiological profiles in a ponderosa pine ecosystem ESA 2003 Annual Meeting, August 3-8, 2003 Savannah, GA

Norris TB, Wraith JM, Castenholz RW and McDermott TR (2002) Soil Microbial Community Structure across a Thermal Gradient following a Geothermal Heating Event. *Applied and Environmental Microbiology* 68:6300-6309

Pandit A and Maheshwari R (1996) Life-history of *Neurospora intermedia* in a sugar cane field. *Journal of Biosciences* 21:57-79

Pietikainen J, Hiukka R and Fritze H (2000) Does short-term heating of forest humus change its properties as a substrate for microbes? *Soil Biology and Biochemistry* 32:277-288

- Prieto-Fernández A, Acea MJ and Carballas T (1998) Soil microbial and extractable C and N after wildfire. *Biology and Fertility of Soils* 27:132-142
- Raison RJ (1979) Modification of the soil environment by vegetation fires, with particular reference to nitrogen transformations: a review. *Plant Soil* 51:73-108
- Renkin RA and Despain DG (1992) Fuel moisture, forest type and lightning caused fire in Yellowstone National Park. *Canadian Journal of Forest Research* 22:37-45
- Ross DJ, Speir TW, Tate KR and Feltham CW (1997) Burning in a New Zealand snow-tussock grassland: Effects on soil microbial biomass and nitrogen and phosphorous availability. *New Zealand Journal of Ecology* 21:63-71
- Sharpley A (2000) Phosphorous availability. In: Sumner, M.E., (ed. *Handbook of soil science*. CRC, Boca Raton, Fl., ppD18-D38
- Stracher GB (1995) The Anthracite Smokers of Eastern Pennsylvania: P_{S2}-T Stability Diagram by TL Analysis. *Mathematical Geology* 27:499-511
- Tobin-Janzen T, Shade A, Marshall L, Torres K, Beblo C, Janzen C, Lenig J, Martinez A and Ressler D (2005) Nitrogen Changes and Domain Bacteria Ribotype Diversity in Soils Overlying the Centralia Pennsylvania Underground Coal Mine Fire. *Soil Science* 170:1-10
- Treseder KK, Mack MC and Cross A (2004) Relationships among fires, fungi, and soil dynamics in Alaskan boreal forests. *Ecological Applications* 14:1826-1838
- Trifonoff KM (2000) The mine fire in Centralia, Pennsylvania. *Pennsylvania Geographer* 38:3-24
- UNECE, FAO, Organization IL and Center GFM (2000) Baltic Exercise for Fire Information and Resources Exchange -BALTEX FIRE 2000, 5-9 June 2000 Kuopio, Finland. Conference Report. Helsinki
- UNECE and FAO (2001) Forest fire statistics 1998-2000. *Timber Bulletin LIV ECE/TIM/BULL/54/4*. Rome, FAO
- Vazquez FJ, Acea MJ and Carballas T (1993) Soil microbial populations after wildfire. *FEMS Microbiology Ecology* 13:93-103
- Westerling AL, Hidalgo HG, Cayan DR and Swetnam TW (2006) Warming and Earlier Spring Increase Western U.S. Forest Wildfire Activity. *Science* 313:940-943

Weston CJ and Attiwill PM (1996) Clearfelling and burning effects on nitrogen mineralization and leaching in soils of old-age *Eucalyptus regnans* forests. *Forest Ecology and Management* 89:13-24

Yeager CM, Northup DE, Grow CC, Barns SM and Kuske CR (2005) Changes in Nitrogen-Fixing and Amonia-Oxidizing Bacterial Communities in Soil. *Applied Envrionmental Microbiology* 71:2713-2722

Zhou F, Ren W, Wang D, Song T, Li X and Zhang Y (2006) Application of three-phase foam to fight an extraordinarily serious coal mine fire. *International Journal of Coal Geology* 67:95-100